

# CROSSCAP NUMBERS OF 2-BRIDGE KNOTS

MIKAMI HIRASAWA AND MASAKAZU TERAGAITO

ABSTRACT. We present a practical algorithm to determine the minimal genus of non-orientable spanning surfaces for 2-bridge knots, called the crosscap numbers. We will exhibit a table of crosscap numbers of 2-bridge knots up to 12 crossings (all 362 of them).

## 1. INTRODUCTION

For a knot  $K$  in the 3-sphere  $S^3$ , there is a connected compact embedded surface  $F$  in  $S^3$  whose boundary is  $K$ . In particular,  $F$  can be chosen to be orientable, and then it is called a Seifert surface for  $K$ . The genus  $g(K)$  of  $K$  is the minimal number of genera of all Seifert surfaces for  $K$ . Thus the unknot is the only knot of genus zero.

On the other hand, we can choose the above  $F$  to be non-orientable, for example, by adding a half-twisted band to a Seifert surface. In this paper, such  $F$  is referred to as a *non-orientable spanning surface* for  $K$ . We define the *crosscap number*  $\gamma(K)$  of a non-trivial knot  $K$  as the minimal number of the first betti numbers  $\beta_1$  of all non-orientable spanning surfaces for  $K$ , and set  $\gamma(\text{unknot}) = 0$  for convenience. We call  $\gamma(K)$  the crosscap number because it counts the number of ‘crosscap summands’ in the closed surface obtained by capping off a non-orientable spanning surface with a disk, which is well known to be a connected sum of projective planes. In the literature, a crosscap number is also called a non-orientable genus [9]. For a non-trivial knot  $K$ , if a non-orientable spanning surface  $F$  satisfies  $\beta_1(F) = \gamma(K)$ , then  $F$  is called a *minimal genus non-orientable spanning surface* for  $K$ .

In general, it is very hard to determine the crosscap number for a given knot. Any minimal genus Seifert surface becomes a non-orientable spanning surface for the same knot if we attach a small half-twisted band as above, and hence we have an obvious inequality  $\gamma(K) \leq 2g(K) + 1$ . There are only a few results about crosscap numbers of knots. Clark [4] introduced the notion of crosscap number and pointed out that  $\gamma(K) = 1$  if and only if  $K$  is a 2-cabled knot. He also asked the existence of a knot satisfying the equality  $\gamma(K) = 2g(K) + 1$ , and Murakami and Yasuhara [8] came up with the first example, showing  $\gamma(7_4) = 3$  algebraically. In [11], the crosscap numbers of torus knots are completely determined.

The purpose of this paper is to determine the crosscap numbers of 2-bridge knots, which form a special but important class of knots. For 2-bridge knots, Hatcher and Thurston [7] constructed all incompressible, boundary-incompressible orientable or non-orientable spanning surfaces. However, for the 2-bridge knot  $7_4$ , a minimal genus non-orientable spanning surface can be realized only by a boundary-compressible surface. Then Bessho [1] proved that any incompressible,

---

2000 *Mathematics Subject Classification*. Primary 57M25.

*Key words and phrases*. knot, crosscap number, 2-bridge knot, non-orientable spanning surface.

The first author was partially supported by MEXT, Grant-in-Aid for Young Scientists (B) 14740048.

boundary-compressible spanning surface for a 2-bridge knot becomes an incompressible, boundary-incompressible surface after several boundary-compressions. Therefore, theoretically, we can obtain  $\gamma(K)$  as follows:

For a 2-bridge knot  $K$ , generate all incompressible, boundary-incompressible spanning surfaces according to [7]. Let  $n$  be the minimal first betti number of them. Then if  $n$  is realized by a non-orientable spanning surface, then  $\gamma(K) = n$ , and otherwise  $\gamma(K) = n + 1$ . Here,  $n$  equals the minimal length of all continued fraction expansions for  $K$ .

However, an effective algorithm to determine  $n$  was missing, and one could not tell, for example, for which 2-bridge knots, the equality  $\gamma(K) = 2g(K) + 1$  holds.

In the following, we present a practical algorithm to find a shortest continued fraction expansion for all rational numbers representing a 2-bridge knot  $K$ . This enables us to determine the crosscap number from any continued fraction expansion for  $K$ . The main tool is so-called the modular diagram, whose vertices correspond to rational numbers, on which we introduce the notion of depth. In Section 6, we exhibit a table of crosscap numbers of 2-bridge knots up to 12 crossings (all 362 of them).

## 2. STATEMENT OF RESULTS

Let  $K$  be a 2-bridge knot  $S(q, p)$  in Schubert's notation. Here,  $p$  and  $q$  are coprime integers, and  $q$  is odd. As is well-known,  $S(q, p)$  and  $S(q', p')$  are equivalent if and only if  $q' = q$  and  $p' \equiv p^{\pm 1} \pmod{q}$ , and  $S(q, -p)$  gives the mirror image of  $S(q, p)$ .

Consider a *subtractive* continued fraction expansion of  $p/q$  (see [7])

$$\frac{p}{q} = r + [b_1, b_2, \dots, b_n] = r + \frac{1}{b_1 - \frac{1}{b_2 - \frac{1}{b_3 - \frac{1}{\ddots - \frac{1}{b_n}}}}}$$

where  $r, b_i \in \mathbb{Z}$  and  $b_i \neq 0$ . The *length* of this expansion is  $n$ . Then  $K$  is the boundary of the surface obtained by plumbing  $n$  bands in a row, the  $i$ th band having  $b_i$  half-twists (right-handed if  $b_i > 0$  and left-handed if  $b_i < 0$ ). If some  $b_i$  is odd, then the expansion is said to be of *odd type*. Otherwise, it is of *even type*. Any fraction has expansions of odd type and even type, e.g.,  $1/3 = 1 - 2/3 = 1 + [-2, -2] = 1 + [-1, 2]$ . In this paper, an expansion always means a subtractive one. We remark the following equality:

$$r + [a_1, -a_2, a_3, -a_4, \dots, (-1)^{n-1} a_n] = r + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \frac{1}{\ddots + \frac{1}{a_n}}}}}$$

The crosscap number of a 2-bridge knot  $K$  can be described in terms of the length of expansion corresponding to  $K$ . The first theorem is due to Bessho, but we will give its proof for reader's convenience in Section 3.

**Theorem 2.1** (Bessho [1]). *Let  $K$  be a 2-bridge knot.*

- (1) The crosscap number  $\gamma(K)$  equals the minimal length of all expansions of odd type of all fractions corresponding to  $K$ .
- (2) If a minimal genus non-orientable spanning surface  $F$  for  $K$  is boundary-compressible, then  $F$  is obtained from a minimal genus Seifert surface for  $K$  by attaching a Möbius band as in Figure 1.




FIGURE 1. Möbius bands attached to Seifert surfaces

We present a practical algorithm to obtain a shortest expansion from any one of  $p/q$ .

**Theorem 2.2.** *Let  $p/q = r + [b_1, b_2, \dots, b_n]$  be an expansion obtained from an arbitrary expansion of  $p/q$  by fully reducing the length by a repetition of the following three reductions. Then  $n$  is the minimal length of all expansions of  $p/q$ .*

- (1) Removal of coefficient 0.

$$\begin{aligned} [\dots, a, 0, b, \dots] &= [\dots, a + b, \dots], \\ [\dots, a, b, 0] &= [\dots, a], \\ [0, a, b, \dots] &= -a + [b, \dots]. \end{aligned}$$

- (2) Removal of coefficient  $\varepsilon = \pm 1$ .

$$\begin{aligned} [\dots, a, \varepsilon, b, \dots] &= [\dots, a - \varepsilon, b - \varepsilon, \dots], \\ [\dots, a, \varepsilon] &= [\dots, a - \varepsilon], \\ r + [\varepsilon, a, \dots] &= (r + \varepsilon) + [a - \varepsilon, \dots]. \end{aligned}$$

- (3) Removal of a subsequence  $2\varepsilon, 2\varepsilon$  or  $2\varepsilon, 3\varepsilon, \dots, 3\varepsilon, 2\varepsilon$ . (Here,  $\varepsilon = \pm 1$ , and possibly  $m = 2$ .)

$$\begin{aligned} [\dots, a, \underbrace{2\varepsilon, 3\varepsilon, \dots, 3\varepsilon, 2\varepsilon}_m, b, \dots] &= [\dots, a - \varepsilon, \underbrace{-3\varepsilon, -3\varepsilon, \dots, -3\varepsilon}_{m-1}, b - \varepsilon, \dots], \\ [\dots, a, \underbrace{2\varepsilon, 3\varepsilon, \dots, 3\varepsilon, 2\varepsilon}_m] &= [\dots, a - \varepsilon, \underbrace{-3\varepsilon, -3\varepsilon, \dots, -3\varepsilon}_{m-1}], \\ r + [\underbrace{2\varepsilon, 3\varepsilon, \dots, 3\varepsilon, 2\varepsilon}_m, a, \dots] &= (r + \varepsilon) + [\underbrace{-3\varepsilon, -3\varepsilon, \dots, -3\varepsilon}_{m-1}, a - \varepsilon, \dots], \\ r + [\underbrace{2\varepsilon, 3\varepsilon, \dots, 3\varepsilon, 2\varepsilon}_m] &= (r + \varepsilon) + [\underbrace{-3\varepsilon, -3\varepsilon, \dots, -3\varepsilon}_{m-1}]. \end{aligned}$$

In Theorem 2.2, we fix a fraction  $p/q$ . Although there are infinitely many fractions corresponding to a 2-bridge knot, the next theorem guarantees that we can start from any fraction.

**Theorem 2.3.** *Let  $K$  be a 2-bridge knot. If the reduction in Theorem 2.2 yields a length  $n$  expansion, then  $n$  is the minimal length of all expansions of all fractions corresponding to  $K$ .*

The next theorem is the key to determine whether a fraction  $p/q$  admits a shortest expansion of odd type.

**Theorem 2.4.** *Two shortest expansions for  $p/q$  are deformed to each other by a finite repetition of the following, where  $\varepsilon = \pm 1$ :*

$$\begin{aligned} [\dots, a, 2\varepsilon, b, \dots] &= [\dots, a - \varepsilon, -2\varepsilon, b - \varepsilon, \dots], \\ [\dots, a, 2\varepsilon] &= [\dots, a - \varepsilon, -2\varepsilon], \\ r + [2\varepsilon, a, \dots] &= (r + \varepsilon) + [-2\varepsilon, a - \varepsilon, \dots]. \end{aligned}$$

**Theorem 2.5.** *Let  $K = S(q, p)$  be a 2-bridge knot. If a shortest expansion of  $p/q$  obtained by Theorem 2.2 contains an odd coefficient or  $\pm 2$ , then  $\gamma(K) = n$ , otherwise  $\gamma(K) = n + 1$ , where  $n$  is the length of the expansion.*

Remark that if there is a coefficient  $\pm 2$  in an expansion with only even coefficients, then we can apply Theorem 2.4 to obtain an expansion of odd type.

**Example 2.6.** Let  $K = 6_1$  in the knot table (see [10]). It is the 2-bridge knot  $S(9, 2)$ . Then  $2/9 = [5, 2]$ . Thus  $\gamma(K) = 2$  by Theorem 2.5.

It is known that any 2-bridge knot  $K$  has a unique expansion of even type modulo integer parts, and the length of which equals  $2g(K)$ . As a direct corollary to Theorem 2.5, we can completely characterize those 2-bridge knots satisfying the equality  $\gamma(K) = 2g(K) + 1$ .

**Corollary 2.7.** *For a 2-bridge knot  $K$ , the equality  $\gamma(K) = 2g(K) + 1$  holds if and only if there is no coefficient  $\pm 2$  in the (unique) expansion for  $K$  containing only even coefficients.*

**Example 2.8.** Let  $K = 7_4 = S(15, 4)$ . Note that  $K$  has genus one. Since  $4/15 = [4, 4]$ ,  $\gamma(K) = 2g(K) + 1 = 3$  by Corollary 2.7. More examples will be given in Section 5.

Some minimal genus non-orientable surfaces for 2-bridge knots are boundary-incompressible, but others boundary-compressible, and some 2-bridge knots have several such surfaces. This makes a strong contrast to the case of torus knots, where minimal genus non-orientable spanning surfaces are boundary-incompressible and even unique [11].

By the theorems above, we can characterize 2-bridge knots with boundary-compressible minimal genus non-orientable spanning surfaces. It is unknown whether Corollary 2.10 below generalizes to all knots.

**Theorem 2.9.** *Let  $K = S(q, p)$  be a 2-bridge knot, and  $\mathcal{C}$  the set of shortest expansions for  $p/q$ . Then we have:*

- (1)  $\mathcal{C}$  contains an expansion of odd type if and only if any minimal genus non-orientable spanning surface for  $K$  is boundary-incompressible.
- (2)  $\mathcal{C}$  contains no expansion of odd type if and only if any minimal genus non-orientable spanning surface for  $K$  is boundary-compressible.

The following is immediate from Theorem 2.9.

**Corollary 2.10.** *A 2-bridge knot never has two minimal genus non-orientable spanning surfaces such that one is boundary-incompressible and the other is boundary-compressible.*

In Section 4, we give an algorithm to visualize a minimal genus non-orientable spanning surface for 2-bridge knots.

**Theorem 2.11.** *Any 2-bridge knot  $K$  has a Conway diagram  $D$  such that a minimal genus non-orientable spanning surface for  $K$  is obtained as a checker-board surface on  $D$ .*

**Example 2.12.** We note that such diagrams are not unique and not shortest in general. Let  $K = S(15, 4)$  as in Example 2.8 with  $\gamma(K) = 3$ . Then the Conway diagrams  $[2, 1, 5, -1, 3]$  and  $[4, 1, 1, 1, 4]$  representing  $K$  respectively yield a desired surface as a checker-board surface. It is interesting to confirm Theorem 2.1(2) for these surfaces.

### 3. PROOF OF THEOREM 2.1

Let  $K$  be a 2-bridge knot with a minimal genus non-orientable spanning surface  $F$ . Let  $E(K) = S^3 - \text{Int } N(K)$  be its exterior. Then  $F \cap N(K)$  can be assumed to be a collar neighborhood of  $\partial F$  in  $F$ , and hence we will use the same notation  $F$  for  $F \cap E(K)$ .

**Lemma 3.1.**  *$F$  is incompressible in  $E(K)$ .*

*Proof.* Assume not. Let  $D$  be a compressing disk for  $F$ . Then  $\partial D$  is an orientation-preserving loop on  $F$ . Let  $F'$  be the resulting surface from  $F$  by compressing along  $D$ . Then  $\chi(F') = \chi(F) + 2$ . If  $F'$  is disconnected, then it consists of a closed orientable component  $F_1$  and a non-orientable component  $F_2$  with  $\partial F_2 \neq \emptyset$ . Since  $\beta_1(F_1) + \beta_1(F_2) = \beta_1(F)$  and  $\beta_1(F_1) > 0$ , we have  $\beta_1(F_2) < \beta_1(F)$ . This contradicts the minimality of  $\beta_1(F)$ . If  $F'$  is connected and non-orientable, then  $\beta_1(F') = \beta_1(F) - 2$ , a contradiction. Hence  $F'$  is connected and orientable. This means that  $F'$  is a Seifert surface for  $K$ . Then adding a small half-twisted band to  $F'$  gives a non-orientable spanning surface  $R$  for  $K$  with  $\beta_1(R) = \beta_1(F') + 1 = \beta_1(F) - 1$ , a contradiction.  $\square$

*Proof of Theorem 2.1.* (1) Let  $p/q = r + [b_1, b_2, \dots, b_n]$  be an expansion of odd type of some fraction  $p/q$  for  $K$ . We assume that the length  $n$  is minimal among all expansions of odd type of all fractions for  $K$ . The surface obtained by plumbing  $n$  bands corresponding to this expansion in the usual way gives a non-orientable spanning surface for  $K$  with the first betti number  $n$ . Thus  $\gamma(K) \leq n$ .

The argument to show  $n \leq \gamma(K)$  is divided into two cases according to the boundary-incompressibility of a minimal genus non-orientable spanning surface  $F$ .

First assume that  $K$  has a minimal genus non-orientable spanning surface  $F$  which is boundary-incompressible. Then it is isotopic to one of the surfaces obtained by plumbing  $k$  bands corresponding to some expansion  $s + [a_1, a_2, \dots, a_k]$  of some fraction for  $K$  with  $s \in \mathbb{Z}$  and  $|a_i| \geq 2$  for each  $i$  by [7, Theorem 1(b)]. Hence  $\gamma(K) = k$ . Since  $F$  is non-orientable, this expansion  $s + [a_1, a_2, \dots, a_k]$  must be of odd type. Thus  $n \leq k$  by the minimality of  $n$ , and hence we have  $n \leq \gamma(K)$ .

Next, assume that any minimal genus non-orientable spanning surface  $F$  for  $K$  is boundary-compressible. Let  $D$  be a boundary-compressing disk such that  $\partial D = \alpha \cup \beta$ , where  $D \cap F = \alpha$  is a properly embedded essential arc in  $F$  and  $D \cap \partial E(K) = \beta$ . Then  $\beta$  intersects  $\partial F$  in two points. If these two points have distinct signs (after orienting  $\beta$  and  $\partial F$  suitably), then  $\beta$  and a subarc of  $\partial F$  bound a disk  $\delta$  in  $\partial E(K)$ . Thus  $D \cup \delta$ , pushed away from  $\partial E(K)$  slightly, gives a compressing disk for  $F$ , which contradicts Lemma 3.1. Hence  $\beta$  intersects  $\partial F$  twice in the same direction. Let  $F_1$  be the surface obtained by boundary-compressing  $F$  along  $D$ . From the above observation,  $F_1$  is a connected surface with connected boundary. Also, we see  $\beta_1(F_1) = \beta_1(F) - 1$ .

**Claim 3.2.**  *$F_1$  is incompressible in  $E(K)$ .*

*Proof.* Let  $N(D) = D \times [-1, 1]$  be a product neighborhood of  $D$  such that  $N(D) \cap F = \partial N(D) \cap F = \alpha \times [-1, 1]$ . Then  $F_1 = (F - N(D) \cap F) \cup (D \times \{-1, 1\})$ . If  $F_1$  is compressible, then it has a compressing disk  $E$  disjoint from  $N(D)$ . Since  $F$

is incompressible,  $\partial E$  bounds a disk  $E'$  in  $F$ . We can choose  $E'$  disjoint from the disk  $\alpha \times [-1, 1]$ . Thus  $\partial E$  bounds a disk in  $F_1$ , a contradiction.  $\square$

If  $F_1$  is orientable, then it is boundary-incompressible. (Any orientable incompressible surface in  $E(K)$  is boundary-incompressible if it has a connected boundary.) If  $F_1$  is non-orientable and boundary-compressible, we continue a boundary-compression. Thus for some  $\ell > 0$  we have a sequence of incompressible surfaces  $F = F_0 \rightarrow F_1 \rightarrow F_2 \rightarrow \cdots \rightarrow F_\ell$  where  $\beta_1(F_i) = \beta_1(F_{i-1}) - 1$  for  $i = 1, 2, \dots, \ell$  and  $F_\ell$  is boundary-incompressible. By [7, Proposition 2],  $\partial F_\ell$  runs once longitudinally on  $\partial E(K)$ .

If  $F_\ell$  is orientable, then it has minimal genus [7, Corollary]. Note that  $\beta_1(F_\ell) = \beta_1(F) - \ell = \gamma(K) - \ell$  and that  $F_\ell$  corresponds to the unique expansion  $r' + [b'_1, b'_2, \dots, b'_m]$  with each  $b'_i$  even. Then  $\beta_1(F_\ell) = m$ . Since  $K$  admits an odd type expansion of length  $m + 1$ ,  $n \leq m + 1 = \beta_1(F_\ell) + 1 = \gamma(K) - \ell + 1 \leq \gamma(K)$ . Thus we have  $n \leq \gamma(K)$ , and so  $\gamma(K) = n$  and  $\ell = 1$ .

If  $F_\ell$  is non-orientable, then  $F_\ell$  is isotopic to some surface obtained by plumbing  $k$  bands and  $n \leq k$  as before. Since  $\beta_1(F_\ell) = k$ ,  $n \leq k = \gamma(K) - \ell < \gamma(K) \leq n$ , a contradiction. Thus such a case never occurs.

(2) If a minimal genus non-orientable spanning surface  $F$  is boundary-compressible, then the above argument shows that boundary-compressing  $F$  gives a minimal genus Seifert surface  $Q$  for  $K$ . Then  $F$  is obtained from  $Q \subset E(K)$  by attaching a band  $b = [0, 1] \times [0, 1] \subset \partial E(K)$  such that  $b \cap Q = [0, 1] \times \{0, 1\}$ . In fact, since  $\partial F$  runs once longitudinally on  $\partial E(K)$ , there are only two possibilities for  $b$  as shown in Figure 1. Thus  $F$  is obtained from  $Q$  by adding a Möbius band locally as desired.  $\square$

#### 4. CALCULATION BY THE MODULAR DIAGRAM

We use the modular diagram  $\mathcal{D}$  as shown in Figure 2 to compute the crosscap numbers of 2-bridge knots. This diagram comes from the action of  $PSL(2, \mathbb{Z})$  on the hyperbolic plane. (But  $\mathcal{D}$  is distorted to space the vertices evenly along the circle.)



The vertices are labelled with  $\mathbb{Q} \cup \{1/0\}$ , inductively: Start with  $1/0$  and  $0/1$  at the ends of the ‘horizontal’ edge. If two vertices of an triangle are already labelled with  $a/b$  and  $c/d$ , then the remaining vertex of the triangle is labelled  $(a+c)/(b+d)$ . (This is the rule to label the vertices on the upper circle only, and for those on the lower circle, regard  $1/0$  and  $0/1$  as  $-1/0$  and  $-0/1$ .)

We call the third vertex the *child* of the first two vertices, which themselves are called the *parents*, and call the edge connecting the parents the *longest side* of a triangle. Note that two vertices  $a/b$  and  $c/d$  are connected by an edge if and only if  $|ad - bc| = 1$ .

We will identify a vertex with the corresponding label for convenience. In fact, all rational numbers appear on the circle with the usual order. That is, if the vertices  $u$  and  $v$  correspond to  $a/b$  and  $c/d$ , respectively, and if  $a/b < c/d$  in  $\mathbb{Q}$ , then  $u$  and  $v$  lie on the circle with the counterclockwise orientation in the order  $u, v$ .

An edge-path from  $1/0$  to  $p/q$  in  $\mathcal{D}$  corresponds uniquely to an expansion  $p/q = r + [b_1, b_2, \dots, b_n]$ , where the partial sums  $p_i/q_i = r + [b_1, b_2, \dots, b_i]$  ( $p_0/q_0 = r$ ) are the successive vertices of the edge-path. At the vertex  $p_{i-1}/q_{i-1}$  the path turns left or right across  $|b_i|$  triangles, left if  $b_i > 0$  and right if  $b_i < 0$ . See Figure 3. If an edge-path corresponds to an expansion of odd type, then the path is also said to be of odd type.

We assign the *depth*  $d(v)$  to each vertex  $v$  on  $\mathcal{D}$ . First, set  $d(v) = 0$  for  $v \in \mathbb{Z} \cup \{1/0\}$ . If a triangle in  $\mathcal{D}$  has vertices  $a/b$ ,  $c/d$  and  $(a+c)/(b+d)$ , then define

FIGURE 2. The modular diagram  $\mathcal{D}$ FIGURE 3. Edge-paths from  $1/0$  to  $p/q$ 

the depth of the child from those of parents by setting

$$d((a+c)/(b+d)) = \min\{d(a/b), d(c/d)\} + 1.$$

Thus all vertices can be assigned the depths. Notice that if two vertices  $u$  and  $v$  are connected by an edge in  $\mathcal{D}$ , then  $|d(u) - d(v)| \leq 1$ . Also there are only three kinds of triangles in  $\mathcal{D}$  as shown in Figure 4, where depths of vertices are indicated, except triangles with vertices  $\{1/0, n, n+1\}$  of depth 0 where  $n \in \mathbb{Z}$ .

**Lemma 4.1.** *Let  $\ell$  be the length of a shortest edge-path from  $1/0$  to  $p/q \neq 1/0$ . Then  $d(p/q) = \ell - 1$ .*

*Proof.* By the definition of depth, there is an edge-path from  $1/0$  to  $p/q$  of length  $d(p/q) + 1$ . Thus  $\ell \leq d(p/q) + 1$ . Conversely, let  $\xi$  be a shortest edge-path from  $1/0$  to  $p/q$ . Recall that  $|d(u) - d(v)| \leq 1$  for any consecutive vertices  $u, v$  on  $\xi$ .




FIGURE 4. Triangles

In particular,  $d(1/0) = d(v_0) = 0$ , where  $v_0$  is the second vertex on  $\xi$ , and hence  $d(p/q) \leq \ell - 1$ . Thus we have  $d(p/q) = \ell - 1$ .  $\square$

The next theorem gives a criterion for a shortest edge-path in terms of depths.

**Theorem 4.2.** *An edge-path  $\xi : 1/0 \rightarrow v_0 \rightarrow v_1 \rightarrow v_2 \rightarrow \cdots \rightarrow v_n = p/q \neq 1/0$  is shortest if and only if  $d(v_i) = i$  for  $i = 0, 1, 2, \dots, n$ .*

*Proof.* Assume that  $\xi$  is shortest. Then  $d(v_n) = n$  by Lemma 4.1. Since the depth can increase by at most one along  $\xi$  and  $d(v_0) = 0$ ,  $d(v_i) = i$  for each  $i$ .

Conversely, since  $d(v_n) = n$ , any edge-path from  $1/0$  to  $p/q$  has length at least  $n + 1$  by Lemma 4.1. Thus we can conclude that  $\xi$  is shortest.  $\square$

*Proof of Theorem 2.2.* Let  $p/q = r + [b_1, b_2, \dots, b_n]$  be an expansion fully reduced by the reductions in the statement of Theorem 2.2. Let  $\xi : 1/0 \rightarrow v_0 \rightarrow v_1 \rightarrow \cdots \rightarrow v_n = p/q$  be the edge-path corresponding to the expansion. Suppose that the expansion is not of minimal length, that is,  $\xi$  is not shortest. Then the sequence  $\mathcal{S}$  of depths  $d(v_0), d(v_1), d(v_2), \dots, d(v_n)$  is not strictly increasing by Theorem 4.2. Notice that  $d(v_0) = 0$ .


First, suppose that  $\mathcal{S}$  contains  $i - 1, i, i - 1$  as a subsequence. Then we can see that there is a triangle of Figure 4(1) such that  $\xi$  runs along the shorter two edges. This means that some coefficient  $b_j$  is  $\pm 1$ , a contradiction.

If  $\mathcal{S}$  contains  $0, 0$ , then  $b_1 = \pm 1$ , a contradiction. Thus  $\mathcal{S}$  contains  $i - 1, i, i, \dots, i$  ( $i$  is repeated  $k (\geq 2)$  times) for some  $i \geq 1$ . We choose  $i$  minimal among such subsequences of  $\mathcal{S}$ .


Let  $u_1$  and  $u_2$  be the depth  $i$  vertices on  $\xi$ , appearing in the order  $u_2, u_1$ . We can suppose that the vertex before  $u_2$  on  $\xi$  has depth  $i - 1$ . There is the unique triangle  $T_1$  which contains the edge between  $u_1$  and  $u_2$  as one of two shorter edges. Without loss of generality, we can assume that  $T_1$  has the form of Figure 4(3). Let  $w_1$  be the remaining vertex of  $T_1$ . If  $u_2$  is the child of  $\{u_1, w_1\}$ , then  $\xi$  contains the edges  $w_1 \rightarrow u_2 \rightarrow u_1$ . Then some  $b_j = -1$ , a contradiction. Hence  $u_1$  is the child of  $\{u_2, w_1\}$ . Let  $T_2$  be the (unique) triangle sharing the edge between  $u_2$  and  $w_1$  with  $T_1$ , and let  $u_3$  be the remaining vertex of  $T_2$ . See Figure 5(1). (Since  $d(w_1) = i - 1$  and  $d(u_2) = i$ ,  $u_3$  is located in this position.) Then  $d(u_3) = i$  or  $i - 1$ . If  $d(u_3) = i$ , then  $\xi$  contains  $w_1 \rightarrow u_2 \rightarrow u_1$ , so some coefficient is  $-1$ . Thus  $d(u_3) = i - 1$ . If  $\xi$  contains  $w_1 \rightarrow u_2 \rightarrow u_1$ , then some coefficient is  $-1$  again. Hence  $\xi$  contains  $u_3 \rightarrow u_2 \rightarrow u_1$ .

If  $i = 1$ , then  $d(u_3) = d(w_1) = 0$ . By the minimality of  $i$ ,  $u_3 = v_0$ . Thus  $\xi$  contains  $1/0 \rightarrow u_3 \rightarrow u_2 \rightarrow u_1$ , so  $b_1 = b_2 = -2$ , a contradiction. (If  $T_1$  has the form of Figure 4(2), then we encounter 1 or 2, 2.)



FIGURE 5. Paths  $u_2 \rightarrow u_1$  and  $u_3 \rightarrow u_2 \rightarrow u_1$ 

Suppose  $i \geq 2$ . Let  $T_3$  be the triangle sharing the edge between  $u_3$  and  $w_1$  with  $T_2$ , and let  $w_2$  be the remaining vertex of  $T_3$ . If  $u_3$  is the child of  $\{w_1, w_2\}$ , then  $d(w_2) = i - 2$ , and  $\xi$  contains  $w_2 \rightarrow u_3 \rightarrow u_2 \rightarrow u_1$ . Then we have  $-2, -2$  in the coefficients, a contradiction. Hence  $w_1$  is the child of  $\{u_3, w_2\}$ , and  $d(w_2) = i - 2$ . See Figure 5(2). Let  $T_4$  be the triangle sharing the edge between  $u_3$  and  $w_2$  with  $T_3$ , and let  $u_4$  be the remaining vertex of  $T_4$ . Then  $u_3$  is the child of  $\{u_4, w_2\}$ , and so  $d(u_4) = i - 1$  or  $i - 2$ . If  $d(u_4) = i - 1$ , then  $\xi$  contains  $w_2 \rightarrow u_3 \rightarrow u_2 \rightarrow u_1$ . Then we have  $-2, -2$  in the coefficients. Hence  $d(u_4) = i - 2$ . If  $\xi$  contains  $w_2 \rightarrow u_3 \rightarrow u_2 \rightarrow u_1$ , then we have  $-2, -2$  again. Thus  $\xi$  contains  $u_4 \rightarrow u_3 \rightarrow u_2 \rightarrow u_1$ . See Figure 6(1). If  $i = 2$ , then  $d(u_4) = d(w_2) = 0$  and hence  $u_4 = v_0 (\neq 1/0)$ . Thus  $\xi$  contains  $1/0 \rightarrow u_4 \rightarrow u_3 \rightarrow u_2 \rightarrow u_1$ , so  $b_1 = -2$ ,  $b_2 = -3$  and  $b_3 = -2$ , a contradiction. Thus  $i \geq 3$ .

FIGURE 6. Paths  $u_4 \rightarrow u_3 \rightarrow u_2 \rightarrow u_1$  and  $u_5 \rightarrow u_4 \rightarrow u_3 \rightarrow u_2 \rightarrow u_1$

Continuing this process, we obtain the triangles  $T_1, T_2, \dots, T_{2i}$ , where  $T_{2m-1}$  is the form of Figure 4(3), and  $T_{2m}$  is of Figure 4(1). Also,  $T_1$  has vertices  $\{u_1, u_2, w_1\}$ ,  $T_{2m-1}$  ( $m \geq 2$ ) has vertices  $\{w_{m-1}, w_m, u_{m+1}\}$ , and  $T_{2m}$  ( $m \geq 1$ ) has vertices  $\{u_{m+1}, u_{m+2}, w_m\}$ , and  $d(u_1) = i$ ,  $d(u_j) = i - j + 2$  for  $j \geq 2$ ,  $d(w_j) = i - j$  for  $j \geq 1$ . The path  $\xi$  contains the edges  $u_{i+2} \rightarrow u_{i+1} \rightarrow \dots \rightarrow u_2 \rightarrow u_1$ . Since  $d(u_{i+2}) = 0$ ,  $u_{i+2} = v_0$ . Then  $-2, -3, \dots, -3, -2$  appears in the coefficients, a contradiction. (If  $T_1$  has the form of Figure 4(2), then we encounter  $1$  or  $2, 3, 3, \dots, 3, 2$  in the coefficients.)  $\square$

**Lemma 4.3.** *Let  $p/q$  and  $p'/q'$  be two fractions for a 2-bridge knot  $K$ . Then there exists a one-to-one correspondence between the set of all expansions for  $p/q$  and that of  $p'/q'$ , such that the correspondence preserves both length and type of expansions.*

*Proof.* Since both  $p/q$  and  $p'/q'$  represent the same knot,  $q = q'$  and (i)  $p \equiv p' \pmod{q}$  or (ii)  $pp' \equiv 1 \pmod{q}$ . Suppose  $p/q = p'/q + s$ , where  $s \in \mathbb{Z}$ . Then for any expansion  $r + [a_1, a_2, \dots, a_n]$  of  $p/q$ , we can associate  $r - s + [a_1, a_2, \dots, a_n]$  of  $p'/q$ . Therefore, it suffices to establish a one-to-one correspondence only for  $p/q$  and  $p'/q$  lying between 0 and 1. Under such a restriction, suppose that  $pp' \equiv 1 \pmod{q}$ . Then it is well known that if  $[a_1, a_2, \dots, a_n] = p/q$  then  $[a_n, \dots, a_2, a_1] = p'/q$ . Thus we can establish a required one-to-one correspondence.  $\square$

*Proof of Theorem 2.3.* Let  $p/q$  be a fraction corresponding to  $K$ . Let  $n$  be the minimal length of an expansion of  $p/q$  obtained by the reduction of Theorem 2.2. If another fraction for  $K$  admits an expansion of length shorter than  $n$ , then  $p/q$  has an expansion of the same length by Lemma 4.3. This contradicts the minimality of  $n$ .  $\square$

Next, we consider when there exists a shortest edge-path from  $1/0$  to  $p/q$ , which is of odd type. A *rectangle* in  $\mathcal{D}$  is the union of two triangles sharing one edge.

**Lemma 4.4.** *Assume that a shortest edge-path  $\xi$  from  $1/0$  to  $p/q$  is not of odd type. If there is a rectangle in  $\mathcal{D}$  containing two successive edges of  $\xi$ , then there is a shortest edge-path of odd type from  $1/0$  to  $p/q$ .*

*Proof.* Without loss of generality, we can assume that the successive two edges on  $\xi$ ,  $v_{i-2} \rightarrow v_{i-1}$  and  $v_{i-1} \rightarrow v_i$  lie on a rectangle which is the union of two triangles whose vertices  $\{v_{i-2}, v_{i-1}, v'_{i-1}\}$  and  $\{v_{i-1}, v'_{i-1}, v_i\}$ . Then we replace these two edges by  $v_{i-2} \rightarrow v'_{i-1}$  and  $v'_{i-1} \rightarrow v_i$ . This will change  $\xi$  to a new shortest edge-path of odd type. For, if  $v_{i-2} \neq 1/0$ , then the expansion corresponding to  $\xi$  changes from  $r + [\dots, a, \pm 2, b, \dots]$  to  $r + [\dots, a \mp 1, \mp 2, b \mp 1, \dots]$ , or  $r + [\dots, a, \pm 2]$  to  $r + [\dots, a \mp 1, \mp 2]$ . If  $v_{i-2} = 1/0$ , then  $v_{i-1} \in \mathbb{Z}$ . Thus  $\xi$  corresponds to an expansion  $p/q = v_{i-1} + [\pm 2, a, \dots]$ . Then the new edge-path corresponds to  $v_{i-1} \pm 1 + [\mp 2, a \mp 1, \dots]$ .  $\square$

The deformation used in the proof of Lemma 4.4 is referred to as a *rectangle move*.

*Proof of Theorem 2.4.* Let

$$\xi : 1/0 \rightarrow v_0 \rightarrow v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_n = p/q,$$

$$\xi' : 1/0 \rightarrow v'_0 \rightarrow v'_1 \rightarrow v'_2 \rightarrow \dots \rightarrow v'_n = p/q$$

be two shortest edge-paths from  $1/0$  to  $p/q$ . Recall that each of  $v_i$  and  $v'_i$  has depth  $i$  for any  $i$  by Theorem 4.2. In particular,  $v_0, v'_0 \in \mathbb{Z}$ .

Suppose  $v_0 \neq v'_0$ . We may assume that  $v_0 < v'_0$ . Since  $p/q \in (v_0 - 1, v_0 + 1)$  and  $p/q \in (v'_0 - 1, v'_0 + 1)$ ,  $v'_0 - v_0 = 1$ . In fact,  $v_1$  and  $v'_1$  lie in the interval  $(v_0, v'_0)$ . Let  $u$  be the child of  $\{v_0, v'_0\}$ . Then  $d(u) = 1$ . If neither of  $v_1$  nor  $v'_1$  is  $u$ , then we would have  $p/q < u < p/q$ , a contradiction. Hence we may assume  $v'_1 = u$ . Then a

single application of rectangle move on  $\xi'$  changes the edges  $1/0 \rightarrow v'_0 \rightarrow v'_1 = u$  to  $1/0 \rightarrow v_0 \rightarrow u$ .

Suppose that  $v_i = v'_i$  for  $0 \leq i \leq k$  and  $v_{k+1} \neq v'_{k+1}$  for some  $k \geq 0$ . If  $v_{k+1} < v_k = v'_k < v'_{k+1}$  or  $v_{k+1} > v_k = v'_k > v'_{k+1}$ , then we would have  $p/q < v_k = v'_k < p/q$ , a contradiction. Hence we consider only the case where  $v_{k+1}, v'_{k+1} > v_k = v'_k$ . The case where  $v_{k+1}, v'_{k+1} < v_k = v'_k$  is similar. Without loss of generality, assume  $v'_{k+1} < v_{k+1}$ . Then  $p/q$  lies in the interval  $(v'_{k+1}, v_{k+1})$ . If there is a vertex  $u$  with depth  $k+1$  inside the interval, then we have  $p/q < u < p/q$ , a contradiction. Hence there are a triangle  $\Delta_1$  whose vertices are  $v_k, v_{k+1}$  and  $v'_{k+1}$ , and a triangle  $\Delta_2$  whose vertices are  $v_{k+1}, v'_{k+1}$  and  $w$ . Here,  $v_{k+1}, v'_{k+1}$  are the parents of  $w$ . If  $w = p/q$ , then  $v_{k+2} = v'_{k+2}$ . Then  $\xi$  can be changed to  $\xi'$  by the rectangle move on  $\Delta_1 \cup \Delta_2$ . Otherwise  $p/q$  lies in  $(v'_{k+1}, w)$  or  $(w, v_{k+1})$ . Then  $w = v_{k+2}$  in the former case, and  $w = v'_{k+2}$  in the latter. After the rectangle move on  $\Delta_1 \cup \Delta_2$  to  $\xi$  or  $\xi'$ , we have  $v_{k+1} = v'_{k+1}$ . Thus,  $\xi$  can be changed to  $\xi'$  gradually.  $\square$

*Proof of Theorem 2.5.* If a shortest expansion of  $p/q$  obtained by Theorem 2.2 contains an odd coefficient, then  $\gamma(K) = n$  by Theorems 2.1 and 2.3. Let  $\xi$  be the corresponding edge-path and assume that  $\xi$  is of even type. If the expansion contains a coefficient  $\pm 2$ , then there is a rectangle in  $\mathcal{D}$  containing two successive edges of  $\xi$ . Hence a rectangle move creates another shortest edge-path of odd type by Lemma 4.4. Then  $\gamma(K) = n$  as above. Otherwise  $\xi$  is the unique shortest edge-path from  $1/0$  to  $p/q$  by Theorem 2.4.

If another fraction for  $K$  admits an expansion of odd type of length  $n$ , then  $p/q$  also admits an expansion of odd type of length  $n$  by Lemma 4.3. Thus there is no expansion of odd type of length  $n$ , and so  $\gamma(K) = n + 1$ .  $\square$

*Proof of Corollary 2.7.* It is well known that even type expansions for a 2-bridge knot are unique modulo integer parts, and the length equals  $2g(K)$ . If the expansion does not contain  $\pm 2$ , then it is shortest by Theorem 2.2. Hence  $\gamma(K) = 2g(K) + 1$  by Theorem 2.5. If the expansion contains  $\pm 2$ , then there is another expansion of odd type with the same length by Lemma 4.4. This means that  $\gamma(K) \leq 2g(K)$ .  $\square$

*Proof of Theorem 2.9.* Let  $F$  be a minimal genus non-orientable spanning surface for  $K$ . By Lemma 3.1,  $F$  is incompressible.

First, assume that the minimal length  $n$  of expansions of all fractions for  $K$  is realized by an expansion of odd type.

If  $F$  is boundary-compressible, then boundary-compression yields a minimal genus Seifert surface  $S$  for  $K$  with  $\beta_1(S) = \beta_1(F) - 1$  as in the proof of Theorem 2.1. Note that  $S$  is isotopic to a plumbing surface corresponding to the unique expansion with only even coefficients. In particular, such expansion has length  $n - 1$ . This contradicts the minimality of  $n$ . Hence  $F$  is boundary-incompressible. (In this case  $F$  is isotopic to a plumbing surface by [7].)

Next, assume that only expansions of even type realize the minimal length  $n$ . If  $F$  is boundary-incompressible, then  $F$  is isotopic to a plumbing surface which corresponds to some expansion of odd type, which has length  $k \geq n + 1$ . Indeed,  $k = n + 1$  by the minimality of  $\beta_1(F)$ . If  $F$  corresponds to an expansion  $r + [b_1, b_2, \dots, b_{n+1}]$ , then  $|b_i| \geq 2$  for each  $i$  by [7]. This expansion is not shortest, and hence the sequence  $b_1, b_2, \dots, b_{n+1}$  contains a subsequence  $2\varepsilon, 3\varepsilon, \dots, 3\varepsilon, 2\varepsilon$ ,  $\varepsilon = \pm 1$ , where the number of  $3\varepsilon$  may be zero, by Theorem 2.2. But such expansion can be reduced as shown in Theorem 2.2. In particular, we have a shorter expansion of odd type, a contradiction. Thus  $F$  must be boundary-compressible.  $\square$

*Proof of Theorem 2.11.* Let  $K = S(q, p)$  with  $\gamma(K) = \gamma$ . We omit the trivial case  $\gamma = 1$ . Let  $C = [a_1, b_1, a_2, b_2, \dots]$  be a shortest expansion for  $p/q$  among those

of odd type. To be precise take  $[a_1, b_1, \dots, a_{(\gamma+1)/2}]$  if  $\gamma$  is odd and if otherwise, take  $[a_1, b_1, \dots, b_{\gamma/2}]$ . Figure 7(1) represents a Conway diagram for  $K = S(q, p)$  of length  $\gamma$ . Deform it to Figure 7(3) through (2) corresponding to the expansion  $C' = [a_1 - 1, -1, b_1, 1, a_2, -1, b_2, 1, \dots]$ , which ends with  $a_{(\gamma+1)/2} + 1$  (resp. 1) if  $\gamma$  is odd (resp. even). Then we have a desired Conway diagram with a checkerboard surface




FIGURE 7. Deformation of Conway diagram

$F$  with  $\beta(F) = \gamma$ . We remark it is not preferable if  $a_1 - 1 = 0$  or  $a_{(\gamma+1)/2} + 1 = 0$ . In that case, apply the following: Take the mirror image of  $K$  and thus change all the signs of  $C$ , apply the deformation in Figure 7, and then take the mirror image again. Then we obtain a desired Conway diagram for  $K$ . (This modification works since  $C$  has at most one coefficient  $\pm 1$  because of the minimality of  $\gamma$ .)  $\square$

## 5. EXAMPLES

In [12], it is proved that any positive integer can appear as the crosscap number of some pretzel knot. We can show that such examples can be found among 2-bridge knots.

As seen in Section 6, among 2-bridge knots up to 12 crossings, exactly  $7_4$ ,  $8_3$ ,  $9_5$ ,  $10_3$ ,  $11a_{343}$ ,  $11a_{363}$ ,  $12a_{1166}$  and  $12a_{1287}$  do not have a shortest expansion of odd type and hence satisfy the equality  $\gamma(K) = 2g(K) + 1$ . The next example also gives an infinite series of such 2-bridge knots, as a generalization of [8].

**Example 5.1.** Let  $K_{m,n}$  be the 2-bridge knot corresponding to  $[m, 4, 4, \dots, 4]$  of length  $n$  for any  $n \geq 1$ . When  $n = 1$ ,  $K_{m,1} = S(m, 1)$ .

If  $m$  is odd and  $m \geq 3$ , then this expansion is shortest by Theorem 2.2. Thus  $\gamma(K_{m,n}) = n$  by Theorem 2.5. Also, if  $m \neq m'$ , then  $K_{m,n}$  and  $K_{m',n}$  have distinct denominators, and hence they are not equivalent. Thus we have infinitely many 2-bridge knots  $K_{m,n}$  with  $\gamma(K_{m,n}) = n$  for any  $n \geq 1$ .

If  $m$  and  $n$  are even and  $m \geq 4$ ,  $n \geq 2$ , then  $g(K_{m,n}) = n/2$ . By Corollary 2.7,  $\gamma(K_{m,n}) = n + 1$ . Also, distinct  $m$ 's give distinct knots. Thus we have infinitely many 2-bridge knots  $K_{m,n}$  satisfying the equality  $\gamma(K_{m,n}) = 2g(K_{m,n}) + 1$ .

The Murasugi sum of two minimal genus Seifert surfaces gives a minimal genus Seifert surface [6]. Finally, we give the examples of 2-bridge knots, showing that an analogous statement does not hold in non-orientable case. This is a generalization of Bessho's example [1].

**Example 5.2.** For any odd integer  $m \geq 3$ , let  $K_m$  be the 2-bridge knot corresponding to  $[m, 2]$ . Then  $\gamma(K) = 2$  and its minimal genus non-orientable spanning

surface  $F$  is obtained by plumbing two bands with  $m$  and 2 half-twists respectively. Let  $R$  be the Murasugi sum of two copies  $F_1$  and  $F_2$  of  $F$ . Here, the plumbing disks are chosen to lie in the band with 2 half-twists of  $F_1$  and in the band with 2 half-twists of  $F_2$ . Then  $\beta_1(R) = 4$ . But  $\partial R$  is the 2-bridge knot corresponding to  $[m, 2, 2, m]$ . Since  $[m, 2, 2, m] = [m-1, -3, m-1]$ , it has crosscap number 3. Also, distinct  $m$ 's give inequivalent knots. Thus the Murasugi sum of two minimal genus non-orientable spanning surfaces is not necessarily minimal genus.

## 6. TABLE

Here is the table of crosscap numbers of 362 2-bridge knots up to 12 crossings. The numbering of knots with 10 or less crossings follows that of [10]. For 11, 12 crossings knots, we have used Dowker-Thistlethwaite notation. The last column gives a minimal length subtractive continued fraction expansion of  $p/q$ . We chose them to be of odd type except for the ones (indicated by  $*$ ) where the shortest expansion is unique and of even type. We referred to [2] for 2-bridge knots up to 10 crossings, to [3] for those of 11 and 12 crossings, for which we also used a table compiled by David De Wit [5].

| knot             | $p/q$ | $\gamma$ | continued fraction | knot             | $p/q$ | $\gamma$ | continued fraction |
|------------------|-------|----------|--------------------|------------------|-------|----------|--------------------|
| 3 <sub>1</sub>   | 1/3   | 1        | [3]                | 4 <sub>1</sub>   | 2/5   | 2        | [3, 2]             |
| 5 <sub>1</sub>   | 1/5   | 1        | [5]                | 5 <sub>2</sub>   | 3/7   | 2        | [2, -3]            |
| 6 <sub>1</sub>   | 2/9   | 2        | [5, 2]             | 6 <sub>2</sub>   | 4/11  | 2        | [3, 4]             |
| 6 <sub>3</sub>   | 5/13  | 3        | [3, 2, -2]         | 7 <sub>1</sub>   | 1/7   | 1        | [7]                |
| 7 <sub>2</sub>   | 5/11  | 2        | [2, -5]            | 7 <sub>3</sub>   | 4/13  | 2        | [3, -4]            |
| 7 <sub>4</sub>   | 4/15  | 3        | [4, 4]*            | 7 <sub>5</sub>   | 7/17  | 3        | [2, -2, 3]         |
| 7 <sub>6</sub>   | 7/19  | 3        | [3, 3, -2]         | 7 <sub>7</sub>   | 8/21  | 3        | [3, 3, 3]          |
| 8 <sub>1</sub>   | 2/13  | 2        | [7, 2]             | 8 <sub>2</sub>   | 6/17  | 2        | [3, 6]             |
| 8 <sub>3</sub>   | 4/17  | 3        | [4, -4]*           | 8 <sub>4</sub>   | 5/19  | 2        | [4, 5]             |
| 8 <sub>6</sub>   | 10/23 | 3        | [2, -3, 3]         | 8 <sub>7</sub>   | 9/23  | 3        | [3, 2, -4]         |
| 8 <sub>8</sub>   | 9/25  | 3        | [3, 4, -2]         | 8 <sub>9</sub>   | 7/25  | 3        | [4, 2, -3]         |
| 8 <sub>11</sub>  | 10/27 | 3        | [3, 3, -3]         | 8 <sub>12</sub>  | 12/29 | 4        | [3, 2, 4, 2]       |
| 8 <sub>13</sub>  | 11/29 | 3        | [3, 3, 4]          | 8 <sub>14</sub>  | 12/31 | 4        | [3, 2, -2, 2]      |
| 9 <sub>1</sub>   | 1/9   | 1        | [9]                | 9 <sub>2</sub>   | 7/15  | 2        | [2, -7]            |
| 9 <sub>3</sub>   | 6/19  | 2        | [3, -6]            | 9 <sub>4</sub>   | 5/21  | 2        | [4, -5]            |
| 9 <sub>5</sub>   | 6/23  | 3        | [4, 6]*            | 9 <sub>6</sub>   | 5/27  | 3        | [5, -2, 2]         |
| 9 <sub>7</sub>   | 13/29 | 3        | [2, -4, 3]         | 9 <sub>8</sub>   | 11/31 | 3        | [3, 5, -2]         |
| 9 <sub>9</sub>   | 9/31  | 3        | [3, -2, 4]         | 9 <sub>10</sub>  | 10/33 | 3        | [3, -3, 3]         |
| 9 <sub>11</sub>  | 14/33 | 3        | [2, -3, -5]        | 9 <sub>12</sub>  | 13/35 | 3        | [3, 3, -4]         |
| 9 <sub>13</sub>  | 10/37 | 3        | [4, 3, -3]         | 9 <sub>14</sub>  | 14/37 | 3        | [3, 3, 5]          |
| 9 <sub>15</sub>  | 16/39 | 4        | [2, -2, 3, -2]     | 9 <sub>17</sub>  | 14/39 | 3        | [3, 5, 3]          |
| 9 <sub>18</sub>  | 17/41 | 4        | [2, -2, 2, -3]     | 9 <sub>19</sub>  | 16/41 | 4        | [3, 2, -3, 2]      |
| 9 <sub>20</sub>  | 15/41 | 3        | [3, 4, 4]          | 9 <sub>21</sub>  | 18/43 | 4        | [2, -3, -2, 3]     |
| 9 <sub>23</sub>  | 19/45 | 4        | [2, -3, -3, 2]     | 9 <sub>26</sub>  | 18/47 | 4        | [3, 3, 2, -3]      |
| 9 <sub>27</sub>  | 19/49 | 4        | [3, 2, -3, -3]     | 9 <sub>31</sub>  | 21/55 | 4        | [3, 3, 3, 3]       |
| 10 <sub>1</sub>  | 2/17  | 2        | [9, 2]             | 10 <sub>2</sub>  | 8/23  | 2        | [3, 8]             |
| 10 <sub>3</sub>  | 6/25  | 3        | [4, -6]*           | 10 <sub>4</sub>  | 7/27  | 2        | [4, 7]             |
| 10 <sub>5</sub>  | 13/33 | 3        | [3, 2, -6]         | 10 <sub>6</sub>  | 16/37 | 3        | [2, -3, 5]         |
| 10 <sub>7</sub>  | 16/43 | 3        | [3, 3, -5]         | 10 <sub>8</sub>  | 6/29  | 2        | [5, 6]             |
| 10 <sub>9</sub>  | 11/39 | 3        | [4, 2, -5]         | 10 <sub>10</sub> | 17/45 | 3        | [3, 3, 6]          |
| 10 <sub>11</sub> | 13/43 | 3        | [3, -3, 4]         | 10 <sub>12</sub> | 17/47 | 3        | [3, 4, -4]         |
| 10 <sub>13</sub> | 22/53 | 4        | [3, 2, 3, -4]      | 10 <sub>14</sub> | 22/57 | 4        | [3, 2, -2, 4]      |

| knot               | $p/q$  | $\gamma$ | continued fraction | knot               | $p/q$  | $\gamma$ | continued fraction  |
|--------------------|--------|----------|--------------------|--------------------|--------|----------|---------------------|
| 10 <sub>15</sub>   | 19/43  | 3        | [2, -4, -5]        | 10 <sub>16</sub>   | 14/47  | 3        | [3, -3, -5]         |
| 10 <sub>17</sub>   | 9/41   | 3        | [5, 2, -4]         | 10 <sub>18</sub>   | 23/55  | 4        | [2, -3, -2, 4]      |
| 10 <sub>19</sub>   | 14/51  | 3        | [4, 3, 5]          | 10 <sub>20</sub>   | 16/35  | 3        | [2, -5, 3]          |
| 10 <sub>21</sub>   | 16/45  | 3        | [3, 5, -3]         | 10 <sub>22</sub>   | 13/49  | 3        | [4, 4, -3]          |
| 10 <sub>23</sub>   | 23/59  | 4        | [3, 2, -3, 3]      | 10 <sub>24</sub>   | 24/55  | 4        | [2, -3, 2, -3]      |
| 10 <sub>25</sub>   | 24/65  | 4        | [3, 3, -2, 3]      | 10 <sub>26</sub>   | 17/61  | 4        | [4, 2, -2, 3]       |
| 10 <sub>27</sub>   | 27/71  | 4        | [3, 3, 3, -3]      | 10 <sub>28</sub>   | 19/53  | 3        | [3, 5, 4]           |
| 10 <sub>29</sub>   | 26/63  | 4        | [2, -2, 3, 4]      | 10 <sub>30</sub>   | 26/67  | 4        | [3, 2, -3, -4]      |
| 10 <sub>31</sub>   | 25/57  | 4        | [2, -4, -2, 3]     | 10 <sub>32</sub>   | 29/69  | 4        | [2, -3, -3, -4]     |
| 10 <sub>33</sub>   | 18/65  | 4        | [4, 3, 2, -3]      | 10 <sub>34</sub>   | 13/37  | 3        | [3, 6, -2]          |
| 10 <sub>35</sub>   | 20/49  | 4        | [3, 2, 6, 2]       | 10 <sub>36</sub>   | 20/51  | 4        | [3, 2, -4, 2]       |
| 10 <sub>37</sub>   | 23/53  | 4        | [2, -3, 3, -2]     | 10 <sub>38</sub>   | 25/59  | 4        | [2, -3, -4, 2]      |
| 10 <sub>39</sub>   | 22/61  | 4        | [3, 4, -2, 2]      | 10 <sub>40</sub>   | 29/75  | 5        | [3, 2, -2, 2, -2]   |
| 10 <sub>41</sub>   | 26/71  | 4        | [3, 4, 3, -2]      | 10 <sub>42</sub>   | 31/81  | 5        | [3, 3, 2, -2, 2]    |
| 10 <sub>43</sub>   | 27/73  | 4        | [3, 3, -3, -3]     | 10 <sub>44</sub>   | 30/79  | 4        | [3, 3, 4, 3]        |
| 10 <sub>45</sub>   | 34/89  | 5        | [3, 3, 3, 2, -2]   | 11a <sub>13</sub>  | 28/61  | 4        | [2, -6, -3, -2]     |
| 11a <sub>59</sub>  | 20/43  | 3        | [2, -7, -3]        | 11a <sub>65</sub>  | 27/59  | 4        | [2, -5, 2, -2]      |
| 11a <sub>75</sub>  | 36/83  | 4        | [2, -3, 4, 3]      | 11a <sub>77</sub>  | 55/131 | 5        | [2, -3, -3, -3, -3] |
| 11a <sub>84</sub>  | 44/101 | 5        | [2, -3, 3, 2, -2]  | 11a <sub>85</sub>  | 47/107 | 5        | [2, -4, -2, 2, 3]   |
| 11a <sub>89</sub>  | 44/119 | 5        | [3, 3, -3, -2, 2]  | 11a <sub>90</sub>  | 23/87  | 4        | [4, 4, -2, -3]      |
| 11a <sub>91</sub>  | 50/129 | 5        | [3, 2, -3, -3, -3] | 11a <sub>93</sub>  | 41/93  | 4        | [2, -4, -5, -3]     |
| 11a <sub>95</sub>  | 33/73  | 4        | [2, -5, -3, 2]     | 11a <sub>96</sub>  | 50/121 | 5        | [2, -2, 3, 3, 3]    |
| 11a <sub>98</sub>  | 18/77  | 4        | [4, -4, -3, -2]    | 11a <sub>110</sub> | 35/97  | 4        | [3, 4, -3, -3]      |
| 11a <sub>111</sub> | 37/103 | 4        | [3, 5, 3, 3]       | 11a <sub>117</sub> | 49/117 | 5        | [2, -3, -2, 3, 3]   |
| 11a <sub>119</sub> | 34/77  | 4        | [2, -4, -5, -2]    | 11a <sub>120</sub> | 45/109 | 5        | [2, -2, 3, 3, -2]   |
| 11a <sub>121</sub> | 50/119 | 5        | [2, -3, -3, -3, 2] | 11a <sub>140</sub> | 17/65  | 3        | [4, 6, 3]           |
| 11a <sub>144</sub> | 17/73  | 4        | [4, -3, 2, -2]     | 11a <sub>145</sub> | 22/83  | 4        | [4, 4, -3, -2]      |
| 11a <sub>154</sub> | 30/67  | 4        | [2, -4, 3, -2]     | 11a <sub>159</sub> | 46/111 | 5        | [2, -2, 2, -3, -3]  |
| 11a <sub>166</sub> | 14/59  | 3        | [4, -5, -3]        | 11a <sub>174</sub> | 28/79  | 4        | [3, 5, -2, -3]      |
| 11a <sub>175</sub> | 41/105 | 5        | [3, 2, -4, -2, 2]  | 11a <sub>176</sub> | 31/111 | 5        | [4, 2, -3, -2, 2]   |
| 11a <sub>177</sub> | 21/97  | 4        | [5, 3, 3, 3]       | 11a <sub>178</sub> | 34/123 | 5        | [4, 2, -2, -3, -3]  |
| 11a <sub>179</sub> | 20/57  | 3        | [3, 7, 3]          | 11a <sub>180</sub> | 25/89  | 4        | [4, 2, -4, -3]      |
| 11a <sub>182</sub> | 13/73  | 4        | [6, 2, -2, -3]     | 11a <sub>183</sub> | 34/115 | 5        | [3, -3, -2, 2, 3]   |
| 11a <sub>184</sub> | 19/87  | 4        | [5, 2, -3, -3]     | 11a <sub>185</sub> | 30/109 | 4        | [4, 3, 4, 3]        |
| 11a <sub>186</sub> | 39/95  | 5        | [2, -2, 3, -2, 2]  | 11a <sub>188</sub> | 14/67  | 3        | [5, 5, 3]           |
| 11a <sub>190</sub> | 18/85  | 4        | [5, 3, -2, -3]     | 11a <sub>191</sub> | 19/83  | 4        | [4, -3, -3, 2]      |
| 11a <sub>192</sub> | 26/97  | 4        | [4, 4, 3, -2]      | 11a <sub>193</sub> | 29/95  | 4        | [3, -4, -3, -3]     |
| 11a <sub>195</sub> | 8/53   | 3        | [7, 3, 3]          | 11a <sub>203</sub> | 11/63  | 3        | [6, 4, 3]           |
| 11a <sub>204</sub> | 30/101 | 4        | [3, -3, -4, -3]    | 11a <sub>205</sub> | 25/91  | 4        | [4, 3, -4, -2]      |
| 11a <sub>206</sub> | 7/47   | 3        | [7, 3, -2]         | 11a <sub>207</sub> | 26/85  | 4        | [3, -4, -3, 2]      |
| 11a <sub>208</sub> | 31/105 | 5        | [3, -3, -2, 2, -2] | 11a <sub>210</sub> | 16/73  | 4        | [5, 2, -3, 2]       |
| 11a <sub>211</sub> | 12/67  | 4        | [6, 2, -3, -2]     | 11a <sub>220</sub> | 23/85  | 4        | [4, 3, -3, 2]       |
| 11a <sub>224</sub> | 27/89  | 4        | [3, -3, 3, 3]      | 11a <sub>225</sub> | 11/53  | 3        | [5, 5, -2]          |
| 11a <sub>226</sub> | 20/71  | 4        | [4, 2, -5, -2]     | 11a <sub>229</sub> | 16/71  | 4        | [4, -2, 3, -2]      |
| 11a <sub>230</sub> | 8/51   | 3        | [6, -3, -3]        | 11a <sub>234</sub> | 5/37   | 3        | [7, -2, 2]          |
| 11a <sub>235</sub> | 22/71  | 4        | [3, -4, 2, -2]     | 11a <sub>236</sub> | 29/99  | 5        | [3, -2, 2, -2, 2]   |
| 11a <sub>238</sub> | 12/65  | 4        | [5, -2, 2, -2]     | 11a <sub>242</sub> | 9/47   | 3        | [5, -4, 2]          |
| 11a <sub>243</sub> | 20/69  | 4        | [3, -2, 4, -2]     | 11a <sub>246</sub> | 13/41  | 3        | [3, -6, 2]          |
| 11a <sub>247</sub> | 2/19   | 2        | [9, -2]            | 11a <sub>306</sub> | 29/105 | 4        | [4, 3, 3, 4]        |
| 11a <sub>307</sub> | 18/83  | 4        | [5, 2, -2, -4]     | 11a <sub>308</sub> | 15/71  | 3        | [5, 4, 4]           |
| 11a <sub>309</sub> | 25/93  | 4        | [4, 3, -2, -4]     | 11a <sub>310</sub> | 14/61  | 3        | [4, -3, -5]         |

| knot               | $p/q$  | $\gamma$ | continued fraction    | knot               | $p/q$  | $\gamma$ | continued fraction   |
|--------------------|--------|----------|-----------------------|--------------------|--------|----------|----------------------|
| 11a <sub>311</sub> | 18/79  | 4        | [4, -3, -2, 3]        | 11a <sub>333</sub> | 14/65  | 3        | [5, 3, 5]            |
| 11a <sub>334</sub> | 9/49   | 3        | [5, -2, 4]            | 11a <sub>335</sub> | 17/75  | 4        | [4, -2, 2, -3]       |
| 11a <sub>336</sub> | 11/59  | 3        | [5, -3, -4]           | 11a <sub>337</sub> | 26/89  | 4        | [3, -2, 3, 4]        |
| 11a <sub>339</sub> | 13/55  | 3        | [4, -4, 3]            | 11a <sub>341</sub> | 19/61  | 3        | [3, -5, -4]          |
| 11a <sub>342</sub> | 4/29   | 2        | [7, -4]               | 11a <sub>343</sub> | 4/31   | 3        | [8, 4] <sup>*</sup>  |
| 11a <sub>355</sub> | 7/45   | 3        | [6, -2, 3]            | 11a <sub>356</sub> | 24/79  | 4        | [3, -3, 2, -3]       |
| 11a <sub>357</sub> | 27/91  | 4        | [3, -3, -3, 3]        | 11a <sub>358</sub> | 5/31   | 2        | [6, -5]              |
| 11a <sub>359</sub> | 10/53  | 3        | [5, -3, 3]            | 11a <sub>360</sub> | 10/57  | 3        | [6, 3, -3]           |
| 11a <sub>363</sub> | 6/35   | 3        | [6, 6] <sup>*</sup>   | 11a <sub>364</sub> | 3/25   | 2        | [8, -3]              |
| 11a <sub>365</sub> | 16/51  | 3        | [3, -5, 3]            | 11a <sub>367</sub> | 1/11   | 1        | [11]                 |
| 12a <sub>38</sub>  | 33/71  | 4        | [2, -6, 2, 3]         | 12a <sub>169</sub> | 23/49  | 3        | [2, -8, -3]          |
| 12a <sub>197</sub> | 32/69  | 4        | [2, -6, 3, 2]         | 12a <sub>204</sub> | 76/173 | 5        | [2, -4, -3, -3, -3]  |
| 12a <sub>206</sub> | 47/105 | 4        | [2, -4, 4, 3]         | 12a <sub>221</sub> | 66/169 | 5        | [3, 2, -4, -3, -3]   |
| 12a <sub>226</sub> | 75/181 | 6        | [2, -2, 2, -2, 2, 3]  | 12a <sub>239</sub> | 40/87  | 4        | [2, -6, -3, 2]       |
| 12a <sub>241</sub> | 57/127 | 5        | [2, -4, 3, 2, -2]     | 12a <sub>243</sub> | 60/133 | 5        | [2, -4, 2, 3, 3]     |
| 12a <sub>247</sub> | 71/163 | 5        | [2, -3, 3, 3, 3]      | 12a <sub>251</sub> | 59/159 | 5        | [3, 3, -3, 2, 3]     |
| 12a <sub>254</sub> | 23/97  | 4        | [4, -4, 2, 3]         | 12a <sub>255</sub> | 28/107 | 4        | [4, 5, -2, -3]       |
| 12a <sub>257</sub> | 80/191 | 6        | [2, -2, 2, 3, -2, -3] | 12a <sub>259</sub> | 52/115 | 4        | [2, -5, -4, -3]      |
| 12a <sub>300</sub> | 68/155 | 5        | [2, -3, 2, 4, 3]      | 12a <sub>302</sub> | 61/147 | 5        | [2, -2, 2, -4, -3]   |
| 12a <sub>303</sub> | 64/153 | 5        | [2, -2, 2, 5, 3]      | 12a <sub>306</sub> | 64/147 | 5        | [2, -3, 3, 3, -2]    |
| 12a <sub>307</sub> | 69/157 | 5        | [2, -4, -3, -3, 2]    | 12a <sub>330</sub> | 43/95  | 4        | [2, -5, -4, 2]       |
| 12a <sub>378</sub> | 45/127 | 4        | [3, 6, 3, 3]          | 12a <sub>379</sub> | 17/71  | 3        | [4, -6, -3]          |
| 12a <sub>380</sub> | 20/77  | 3        | [4, 7, 3]             | 12a <sub>384</sub> | 62/151 | 5        | [2, -2, 3, -3, -3]   |
| 12a <sub>385</sub> | 66/161 | 5        | [2, -2, 4, 3, 3]      | 12a <sub>406</sub> | 74/179 | 6        | [2, -2, 3, 2, -2, 2] |
| 12a <sub>425</sub> | 37/81  | 4        | [2, -5, 3, -2]        | 12a <sub>437</sub> | 65/149 | 5        | [2, -3, 2, -3, -3]   |
| 12a <sub>447</sub> | 43/121 | 4        | [3, 5, -3, -3]        | 12a <sub>454</sub> | 27/103 | 4        | [4, 5, -2, 2]        |
| 12a <sub>471</sub> | 38/85  | 4        | [2, -4, 5, 2]         | 12a <sub>477</sub> | 70/169 | 6        | [2, -2, 2, -2, 3, 2] |
| 12a <sub>482</sub> | 22/93  | 4        | [4, -4, 3, 2]         | 12a <sub>497</sub> | 81/209 | 6        | [3, 2, -3, -2, 2, 3] |
| 12a <sub>498</sub> | 76/207 | 5        | [3, 4, 3, 3, 3]       | 12a <sub>499</sub> | 89/233 | 6        | [3, 3, 3, 3, 2, -2]  |
| 12a <sub>500</sub> | 60/167 | 5        | [3, 4, -2, -3, -3]    | 12a <sub>501</sub> | 55/199 | 5        | [4, 3, 3, 3, 3]      |
| 12a <sub>502</sub> | 37/91  | 4        | [2, -2, 6, 3]         | 12a <sub>506</sub> | 68/185 | 5        | [3, 3, -2, -4, -3]   |
| 12a <sub>508</sub> | 56/129 | 5        | [2, -3, 3, -2, 2]     | 12a <sub>510</sub> | 81/193 | 6        | [2, -2, 2, 3, 3, -2] |
| 12a <sub>511</sub> | 51/125 | 5        | [2, -2, 5, 2, -2]     | 12a <sub>512</sub> | 64/151 | 5        | [2, -3, -4, 2, 3]    |
| 12a <sub>514</sub> | 79/187 | 5        | [2, -3, -4, -3, -3]   | 12a <sub>517</sub> | 52/145 | 4        | [3, 5, 4, 3]         |
| 12a <sub>518</sub> | 34/157 | 5        | [5, 2, -2, -3, -3]    | 12a <sub>519</sub> | 25/111 | 4        | [4, -2, 4, 3]        |
| 12a <sub>520</sub> | 36/133 | 4        | [4, 3, -4, -3]        | 12a <sub>521</sub> | 48/113 | 4        | [2, -3, -6, -3]      |
| 12a <sub>522</sub> | 73/173 | 5        | [2, -3, -3, 3, 3]     | 12a <sub>528</sub> | 67/183 | 5        | [3, 4, 4, 2, -2]     |
| 12a <sub>532</sub> | 33/125 | 4        | [4, 5, 3, -2]         | 12a <sub>533</sub> | 31/137 | 5        | [4, -2, 3, 2, -2]    |
| 12a <sub>534</sub> | 44/163 | 5        | [4, 3, -3, -2, 2]     | 12a <sub>535</sub> | 47/175 | 5        | [4, 3, -2, -3, -3]   |
| 12a <sub>536</sub> | 29/137 | 4        | [5, 4, 3, 3]          | 12a <sub>537</sub> | 50/179 | 5        | [4, 2, -3, -3, -3]   |
| 12a <sub>538</sub> | 13/83  | 4        | [6, -2, 2, 3]         | 12a <sub>539</sub> | 44/145 | 5        | [3, -3, 3, 2, -2]    |
| 12a <sub>540</sub> | 49/165 | 5        | [3, -3, -3, 2, 3]     | 12a <sub>541</sub> | 41/153 | 4        | [4, 4, 4, 3]         |
| 12a <sub>545</sub> | 63/143 | 5        | [2, -4, -3, 2, -2]    | 12a <sub>549</sub> | 26/111 | 4        | [4, -4, -3, 2]       |
| 12a <sub>550</sub> | 34/149 | 5        | [4, -2, 2, 3, 3]      | 12a <sub>551</sub> | 18/103 | 4        | [6, 3, -2, -3]       |
| 12a <sub>552</sub> | 30/131 | 4        | [4, -3, -4, -3]       | 12a <sub>579</sub> | 49/177 | 5        | [4, 2, -2, -4, -3]   |
| 12a <sub>580</sub> | 11/69  | 3        | [6, -4, -3]           | 12a <sub>581</sub> | 36/119 | 4        | [3, -3, 4, 3]        |
| 12a <sub>582</sub> | 39/131 | 4        | [3, -3, -5, -3]       | 12a <sub>583</sub> | 45/161 | 5        | [4, 2, -3, -3, 2]    |
| 12a <sub>584</sub> | 31/143 | 5        | [5, 2, -2, -3, 2]     | 12a <sub>585</sub> | 50/181 | 5        | [4, 3, 3, 3, -2]     |
| 12a <sub>595</sub> | 30/139 | 4        | [5, 3, 4, 3]          | 12a <sub>596</sub> | 14/81  | 3        | [6, 5, 3]            |
| 12a <sub>597</sub> | 26/123 | 4        | [5, 4, 3, -2]         | 12a <sub>600</sub> | 25/109 | 4        | [4, -3, -4, 2]       |
| 12a <sub>601</sub> | 34/127 | 4        | [4, 4, 5, 2]          | 12a <sub>643</sub> | 23/99  | 4        | [4, -3, 3, -2]       |

| knot         | $p/q$  | $\gamma$ | continued fraction   | knot         | $p/q$  | $\gamma$ | continued fraction   |
|--------------|--------|----------|----------------------|--------------|--------|----------|----------------------|
| $12a_{644}$  | 30/113 | 4        | $[4, 4, -3, 2]$      | $12a_{649}$  | 27/127 | 4        | $[5, 3, -3, -3]$     |
| $12a_{650}$  | 46/165 | 5        | $[4, 2, -2, 3, 3]$   | $12a_{651}$  | 17/97  | 4        | $[6, 3, -2, 2]$      |
| $12a_{652}$  | 46/155 | 5        | $[3, -3, -3, 2, -2]$ | $12a_{682}$  | 29/107 | 4        | $[4, 3, -4, 2]$      |
| $12a_{684}$  | 41/135 | 5        | $[3, -3, 2, -2, 2]$  | $12a_{690}$  | 20/89  | 4        | $[4, -2, 5, 2]$      |
| $12a_{691}$  | 12/77  | 4        | $[6, -2, 3, 2]$      | $12a_{713}$  | 39/139 | 5        | $[4, 2, -3, 2, -2]$  |
| $12a_{714}$  | 19/107 | 4        | $[6, 3, 3, -2]$      | $12a_{715}$  | 50/169 | 5        | $[3, -3, -3, -3, 2]$ |
| $12a_{716}$  | 5/43   | 3        | $[9, 2, -2]$         | $12a_{717}$  | 28/89  | 4        | $[3, -6, -2, 2]$     |
| $12a_{718}$  | 41/141 | 5        | $[3, -2, 4, 2, -2]$  | $12a_{720}$  | 21/113 | 4        | $[5, -3, -3, -3]$    |
| $12a_{721}$  | 50/171 | 5        | $[3, -2, 3, 3, 3]$   | $12a_{722}$  | 3/29   | 2        | $[10, 3]$            |
| $12a_{723}$  | 20/63  | 3        | $[3, -7, -3]$        | $12a_{724}$  | 31/107 | 4        | $[3, -2, 5, 3]$      |
| $12a_{726}$  | 19/103 | 4        | $[5, -2, 3, 3]$      | $12a_{727}$  | 46/157 | 5        | $[3, -2, 2, -3, -3]$ |
| $12a_{728}$  | 29/133 | 5        | $[5, 2, -2, 2, -2]$  | $12a_{729}$  | 46/167 | 5        | $[4, 3, 3, -2, 2]$   |
| $12a_{731}$  | 22/105 | 4        | $[5, 4, -2, 2]$      | $12a_{732}$  | 18/95  | 4        | $[5, -3, 2, 3]$      |
| $12a_{733}$  | 14/73  | 3        | $[5, -5, -3]$        | $12a_{736}$  | 43/141 | 5        | $[3, -3, 2, 3, -2]$  |
| $12a_{738}$  | 37/119 | 4        | $[3, -5, -3, -3]$    | $12a_{740}$  | 35/113 | 4        | $[3, -4, 3, 3]$      |
| $12a_{743}$  | 12/79  | 4        | $[7, 2, -2, 2]$      | $12a_{744}$  | 8/61   | 3        | $[8, 3, 3]$          |
| $12a_{745}$  | 8/59   | 3        | $[7, -3, -3]$        | $12a_{758}$  | 31/113 | 4        | $[4, 3, 5, -2]$      |
| $12a_{759}$  | 9/61   | 3        | $[7, 4, -2]$         | $12a_{760}$  | 34/111 | 4        | $[3, -4, -4, 2]$     |
| $12a_{761}$  | 41/139 | 5        | $[3, -2, 2, 4, -2]$  | $12a_{762}$  | 7/51   | 3        | $[7, -3, 2]$         |
| $12a_{763}$  | 30/97  | 4        | $[3, -4, 3, -2]$     | $12a_{764}$  | 39/133 | 5        | $[3, -2, 2, -3, 2]$  |
| $12a_{773}$  | 20/91  | 4        | $[4, -2, -5, 2]$     | $12a_{774}$  | 16/89  | 4        | $[5, -2, -4, 2]$     |
| $12a_{775}$  | 16/87  | 4        | $[5, -2, 3, -2]$     | $12a_{791}$  | 13/63  | 3        | $[5, 6, -2]$         |
| $12a_{792}$  | 24/85  | 4        | $[4, 2, -5, 2]$      | $12a_{796}$  | 11/57  | 3        | $[5, -5, 2]$         |
| $12a_{797}$  | 24/83  | 4        | $[3, -2, 5, -2]$     | $12a_{802}$  | 15/47  | 3        | $[3, -7, 2]$         |
| $12a_{803}$  | 2/21   | 2        | $[11, 2]$            | $12a_{1023}$ | 29/127 | 4        | $[4, -3, -3, -4]$    |
| $12a_{1024}$ | 40/149 | 4        | $[4, 4, 3, 4]$       | $12a_{1029}$ | 19/81  | 3        | $[4, -4, -5]$        |
| $12a_{1030}$ | 19/91  | 3        | $[5, 5, 4]$          | $12a_{1033}$ | 25/107 | 4        | $[4, -3, 2, 4]$      |
| $12a_{1034}$ | 32/121 | 4        | $[4, 5, 2, -3]$      | $12a_{1039}$ | 37/137 | 4        | $[4, 3, -3, -4]$     |
| $12a_{1040}$ | 26/115 | 4        | $[4, -2, 3, 4]$      | $12a_{1125}$ | 23/101 | 4        | $[4, -2, 2, 5]$      |
| $12a_{1126}$ | 26/119 | 4        | $[5, 2, -3, -4]$     | $12a_{1127}$ | 22/97  | 4        | $[5, 2, 3, -4]$      |
| $12a_{1128}$ | 9/59   | 3        | $[7, 2, -4]$         | $12a_{1129}$ | 23/105 | 4        | $[4, -2, -4, 3]$     |
| $12a_{1130}$ | 27/125 | 4        | $[5, 3, 3, -3]$      | $12a_{1131}$ | 11/73  | 3        | $[7, 3, 4]$          |
| $12a_{1132}$ | 40/131 | 4        | $[3, -4, -3, -4]$    | $12a_{1133}$ | 47/159 | 5        | $[3, -2, 2, 3, 4]$   |
| $12a_{1134}$ | 7/53   | 3        | $[8, 2, -3]$         | $12a_{1135}$ | 32/103 | 4        | $[3, -4, 2, 4]$      |
| $12a_{1136}$ | 43/147 | 5        | $[3, -2, 3, 2, -3]$  | $12a_{1138}$ | 14/79  | 3        | $[6, 3, 5]$          |
| $12a_{1139}$ | 18/101 | 4        | $[6, 3, 2, -3]$      | $12a_{1140}$ | 18/97  | 4        | $[5, -2, 2, 4]$      |
| $12a_{1145}$ | 15/79  | 3        | $[5, -4, -4]$        | $12a_{1146}$ | 34/117 | 4        | $[3, -2, 4, 4]$      |
| $12a_{1148}$ | 23/73  | 3        | $[3, -6, -4]$        | $12a_{1149}$ | 4/35   | 2        | $[9, 4]$             |
| $12a_{1157}$ | 5/39   | 2        | $[8, 5]$             | $12a_{1158}$ | 16/77  | 3        | $[5, 5, -3]$         |
| $12a_{1159}$ | 24/113 | 4        | $[5, 3, -2, 3]$      | $12a_{1161}$ | 14/75  | 3        | $[5, -3, -5]$        |
| $12a_{1162}$ | 13/69  | 3        | $[5, -3, 4]$         | $12a_{1163}$ | 24/103 | 4        | $[4, -3, 2, -3]$     |
| $12a_{1165}$ | 16/67  | 3        | $[4, -5, 3]$         | $12a_{1166}$ | 4/33   | 3        | $[8, -4]^*$          |
| $12a_{1273}$ | 11/61  | 3        | $[6, 2, -5]$         | $12a_{1274}$ | 17/95  | 4        | $[6, 2, -2, 3]$      |
| $12a_{1275}$ | 44/149 | 5        | $[3, -3, -2, 2, -3]$ | $12a_{1276}$ | 13/75  | 3        | $[6, 4, -3]$         |
| $12a_{1277}$ | 37/121 | 4        | $[3, -4, -3, 3]$     | $12a_{1278}$ | 6/41   | 2        | $[7, 6]$             |
| $12a_{1279}$ | 10/67  | 3        | $[7, 3, -3]$         | $12a_{1281}$ | 33/109 | 4        | $[3, -3, 3, -3]$     |
| $12a_{1282}$ | 10/63  | 3        | $[6, -3, 3]$         | $12a_{1287}$ | 6/37   | 3        | $[6, -6]^*$          |

## REFERENCES

1. K. Bessho, *Incompressible surfaces bounded by links*, Master thesis (in Japanese), Osaka University, 1994.



2. G. Burde and H. Zieschang, *Knots*, de Gruyter Studies in Mathematics **5**, Walter de Gruyter and Co., Berlin, 1985.
3. J. C. Cha and C. Livingston, *KnotInfo*, an online table of knot invariants at [www.indiana.edu/~knotinfo](http://www.indiana.edu/~knotinfo).
4. B. E. Clark, *Crosscaps and knots*, Internat. J. Math. and Math. Sci. **1** (1978), 113–123.
5. D. De Wit, *The 2-bridge knots of up to 16 crossings*, preprint, [arXiv:math.GT/0409351](https://arxiv.org/abs/math/0409351).
6. D. Gabai, *The Murasugi sum is a natural geometric operation*, In Low-dimensional topology (San Francisco, Calif., 1981), 131–143, Contemp. Math., **20**, Amer. Math. Soc., Providence, RI, 1983.
7. A. Hatcher and W. Thurston, *Incompressible surfaces in 2-bridge knot complements*, Invent. Math. **79** (1985), 225–246.
8. H. Murakami and A. Yasuhara, *Crosscap number of a knot*, Pacific J. Math. **171** (1995), 261–273.
9. ———, *Four-genus and four-dimensional clasp number of a knot*, Proc. Amer. Math. Soc. **128** (2000), 3693–3699.
10. D. Rolfsen, *Knots and links*, Mathematics Lecture Series, **7**, Publish or Perish, Inc., Berkeley, Calif., 1976.
11. M. Teragaito, *Crosscap numbers of torus knots*, Topology Appl. **138** (2004), 219–238.
12. J. Uhing, *Zur Kreuzhaubenzahl von Knoten*, diploma thesis, Universität Dortmund.

DEPARTMENT OF MATHEMATICS, GAKUSHUIN UNIVERSITY, 1-5-1 MEJIRO, TOSHIMA-KU, TOKYO 171-8588, JAPAN

*E-mail address:* [hirasawa@math.gakushuin.ac.jp](mailto:hirasawa@math.gakushuin.ac.jp)

DEPARTMENT OF MATHEMATICS AND MATHEMATICS EDUCATION, FACULTY OF EDUCATION, HIROSHIMA UNIVERSITY, 1-1-1 KAGAMIYAMA, HIGASHI-HIROSHIMA 739-8524, JAPAN

*E-mail address:* [teragai@hiroshima-u.ac.jp](mailto:teragai@hiroshima-u.ac.jp)